



*fast*NLQv2


Application and Implementation in the H1Fitter

The fastNLO Collaboration

Daniel Britzger, Thomas Kluge, Klaus Rabbertz, Fred Stober, Markus Wobisch
(DESY, Liverpool, KIT, KIT, Louisiana Tech University)


H1Fitter user's meeting
12. 12. 2011

- Jet cross sections are very slow to calculate
-> Need of method for **very fast repeated calculation** of cross sections
- **FastNLO** factorizes the cross section calculation for an **a-posteriori** inclusion of pdf's and α_s for e.g. jet-production
- Introduce **set of n discrete** $x_{(i)}$'s
with $x_n < \dots < x_i < \dots < x_0 = 1$
- Around each $x_{(i)}$ define **eigen function** $E^{(i)}(x)$ with:
 $E^{(i)}(x_i) = 1, E^{(i)}(x_j) = 0 (i \neq j), \sum_i E^{(i)}(x) = 1$ for all x
- single pdf is replaced by a linear combination of eigenfunctions
- integrals are replaced by sums
- Better: Usage of **bi-cubic interpolation** and **pdf reweighting**




fastNLO, arXiv:1109:1310v1, 2011

- **Comparison of jet data**
 - STAR @ RHIC
 - H1 and ZEUS @ HERA
 - CDF and D0 @ TeVatron
 - CMS and ATLAS @ LHC
- Data/theory plot
- Compatible with NLO pQCD
- First measurements from LHC



- FastNLO tables come with 3 (4) simultaneous **scale variations tables**
 - e.g. 0.5, 1.0, 2.0 times the nominal scale
- **A posteriori scale variation** of the renormalization scale allows study of **asymmetric scale** variations
 - e.g. 6-points: (1/2,1/2), (1/2,1), (1,1/2), (1,2), (2,1), (2,2)
 - avoiding of rel. 'factor' 4.
- **Improvements in v 2.0**
 - scales get own dimension
 - bicubic interpolation of scale-value to scale nodes
 - typically 6 scale nodes
 - examples already for
 - CMS incl. jets
 - D0 3-jet mass
 - ...






New concept 'multidimensional scale table'



- **Improve FastNLO concept**
 - Store scale **independent** weights
 - Store several 'scale' look-up tables, e.g.:
 - p_t
 - Q
- When evaluating FastNLO cross section:
 - User can **choose every scale composition** from previously stored scales
 - e.g. $\mu_r^2 = (Q^2 + p_t^2) / 2$
 - $\mu_r^2 = Q^2$
 - ...
- Also a-posteriori scale variation for μ_r and μ_f are thus **independently possible** through
 - $\mu_r^2 = c_r \times (Q^2 + p_t^2) / 2$
 - $\mu_r^2 = c_f \times Q^2$
 - ...
- New possibilities for scans of scale dependence
- Examples exist for almost **all DIS** tables
 - accuracy $> 0.005\%$ compared to nlojet++ CS
- pp and ppbar concept proven
 - much larger tables

Cross section bin 15

Zeus Dijets Jets @ High Q^2





More features of FastNLO v2.0



• Technicals

- Automatic scan of smallest x-value
- flexible #x-nodes per bin
- scale gets own dimension
- Arbitrary number of dimensions for binning of observable

• Units

- publication units e.g. pb/GeV
- absolute units [pb]


• Additional contributions

- **Correction factors**
 - non-perturbative corrections
 - with errors
- **Data**
 - including arb. no. of correlated and uncorrelated errors
 - Correlation matrix
- **Electroweak corrections** (calculated?)
- **New physics contributions**
- **Threshold corrections**
 - tables are available
- Normalization options

• Converting tool for v1.4 tables

• New concept of 'multidimensional scale table'

- store multiple scales
- user can **compose** scales a-posteriori from all included variables
- speed-up implemented for 'one-scales' (e.g. in pdf fits)
- automated equidistant x-binning in a function of x



CMS Note-2011-004



Release of v2.0



- **Release still **this month** !!!**
- **Easy installation**
 - autotools
 - No further dependencies (exc. LHAPDF)
 - No ROOT
 - No CERNLIBS
- **C++ and fortran reading tools**
 - both 'universal' for all v2.0 tables





What's on today's door?






FastNLO in H1Fitter





FastNLO in H1Fitter



- FastNLO classes
 - FastNLOReader.cc
 - FastNLOBlockB.cc
 - Alphas.cc
- Wrapper
 - FastNLOInterface.cc
- FastNLOInterface
 - `map<int, FastNLOReader*> gFastNLO_array;`
 - `fastnloinit_(const char *s, const int *idataset, const char *thfile)`
 - `fastnlocalc_(const int *idataset, double *xsec)`
- Data card


```
&Data
  Name = 'CMS inclusive jets'
  Reaction = 'FastNLO jets'

  NDATA = 176
  NColumn = 10
  ColumnType = 'Bin' , 'Bin' , 'Bin' , 'Bin' , 'Bin' , 'Bin' , 'Bin'
, 'Sigma' , 'Error' , 'Error'
  ColumnName = 'EtaBinNumber' , 'pt' , 'ymin' , 'ymax' , 'pt1' , 'pt2' ,
'NPCorr' , 'Sigma' , 'stat' , 'uncor'
  NInfo = 1
  DataInfo = 7000.,
  CInfo = 'sqrt(S)'

  IndexDataset = 77
  TheoryInfoFile = 'theoryfiles/fnl2342b.tab'
  TheoryType = 'FastNLO'
  Percent = True, True
&End
1 19.4 0 0.5 18 21 1.4 1.97e+07 0.6 12.85
```



Technicals



- **FastNLOInterface**

```
fnloreader = new FastNLOReader( thfile );

fnloreader->SetPDFInterface(FastNLOReader::kH1FITTER);
fnloreader->SetAlphasEvolution(FastNLOReader::kQCDNUMInternal);
fnloreader->SetScaleVariation(iscale);

// switching non-pert corr off
fnloreader->SetContributionON(FastNLOReader::kNonPerturbativeCorrection,0,false);
fnloreader->SetContributionON(FastNLOReader::kNonPerturbativeCorrection,1,false);

fnloreader->SetUnits(FastNLOReader:: kPublicationUnits);

gFastNLO_array.insert(pair<int, FastNLOReader*>(*idataset, fnloreader) );
```

- **Features**

- use **consistent alpha_s evolution than QCDNUM** (nf, th. matching, n-loop, MZ)

- **Necessary for input card**






- Set scale information
- CS in publication units or absolute units???

 - H1Fitter rule??

- Threshold corrections ON/OFF



Results HERAPDF1.0 vs. HERAPDF1.0 + CMS incl. jets






HeraPDF1.0 + CMS

HERAPDF1.0 Reference
output.HERAPDF10.ref.NLO.bands



$Q^2 = 4.00 \text{ GeV}^2$

CMS jets impact on PDF experimental uncertainty

perfectly consistent with previous fit by K. Nowak



effect from CMS jets inclusion



Scenario 1
No CMS jets
 $Q^2 = 4.00 \text{ GeV}^2$

Experimental error decreased for high-x gluon








Including D0 incl. jets @ NNLO



D0, arXiv:0802.2400

- using D0 jets
- including threshold corrections $O(\text{nnlo})$



HeraPDF1.0 + D0 (incl. th. corr)

"HERAPDF1.0 @ NNLO"

$Q^2 = 4 \text{ GeV}^2$

$\text{Chi}^2/\text{ndf} = 752.9 / 692$





$\text{Chi}^2/\text{ndf} (\text{D0}) = 75.0 / 110$



Which x-region do we test with jet data?




- E.g. H1 dijets @ high Q^2
 - four bins:
 - low and high Q^2
 - low and high $\langle p_T \rangle$
- Only three contributions in DIS
- Gluon, Delta, Sigma induced processes
- low Q^2 is mostly **gluon induced**
- High Q^2 is mostly **Delta induced**
- 'low' x-region only at low $\langle p_T \rangle$ and low Q^2





Can we do the same for CMS incl. jets?



CMS, PRL 107, 132001, 2011

- CMS inclusive jets
- 176 bins
- 6 rapidity regions

- To which 'x'-regions and to which pdfs are we sensitive to???




Can we do the same for CMS incl. jets?



- In FastNLO we replace x-integrations by sums over n_x -nodes
- Basically $13 \times 13 \times n_x \times n_x$ contributions to the cross section
 - reduced to $7(6) \times n_x \times n_x / 2$ cont. in FastNLO
- Still
 - Cross section can be written for FastNLO like

$$\sigma_{\text{Bin}} = \sum_{f1} \sum_{f2} \sum_{x1} \sum_{x2} (\text{pdf}(f1,x1) \times \text{pdf}(f2,x2) \times A)$$

- with $A(f1,f2,x1,x2,\mu_r,\mu_f) = \sigma_{\text{fnlo}} \times \alpha_s$




Can we do the same for CMS incl. jets?

- In FastNLO we replace x-integrations by sums over n_x -nodes
- Basically $13 \times 13 \times n_x \times n_x$ contributions to the cross section
 - reduced to $7(6) \times n_x \times n_x / 2$ cont. in FastNLO
- Still
 - Cross section can be written for FastNLO like

$$\sigma_{\text{Bin}} = \sum_{f1} \sum_{f2} \sum_{x1} \sum_{x2} (\text{pdf}(f1,x1) \times \text{pdf}(f2,x2) \times A)$$

How much is the contribution to the cross section from flavor i as function of x ???







Can we do the same for CMS incl. jets?



 $f_i | x_{1,i}$

$\times A \times$



$\times A \times$

 $f_i | x_{1,i}$




x-dependent contributions






CMS inclusive jets

Low p_t region
Central rapidity

- 'Integral' of all contributions is $2\sigma_{\text{bin}}$
- negative contributions from cancellations (?)
- This Bin
 - testing PDFs up to 10^{-3}
 - no sensitivity to high x region
 - process mostly gluon dominated
 - but what kind of gluon process?







What kind of contributions?









gluon induced processes are mostly gg processes




for gg: we don't know the 'x' of the other gluon

some gq and qq contribution




Central Rapidity - High pt Jets






significant high-x gg contribution

high-x valence meets medium-x gluon




some medium-x u/d meets supposedly high-x gluon



2.50 < |y| < 3.00
300.0 < pt_{jet} < 468.0 GeV

All flavor types in bin 176



valence and gluon dominated



Conclusion



- **FastNLO**

- v2.0 is (almost) released
- new v2.0 tables become available
- v1.4 tables are converted into 'new' table format
- new concept for multidimensional scale tables

- **FastNLO + H1Fitter**

- C++ version (pre-release) is implemented in H1Fitter
- universal interface to all FastNLO tables
 - pp, ppbar, DIS tables
- Alpha_s evolution is identical with QCDNUM
- NNLO fits with jet data (incl. thr. corr.) is principally possible




- **Todo**

- Some parameters should be implemented in steering
- Cross sections in pb or pb/[BinWidth] in H1Fitter ?

- **Open questions**




- Which scales should be used?
- How to determine 'theory unc.' from scale variations
 - how to do it for multiple jet cross sections (DIS,pp,ppbar?)
- How to treat theory uncertainty?
 - Fit with +/- scale variation -> additional pair of pdf-error-set ?





we do not know the 'x' of the other gluon likely: low-x gluon meets high-x gluon

lowest x contributions



gluon dominated

some valence contributions